

Is there a cure?

Yes. Usually, your doctor will ask you to insert an antifungal cream or a suppository inside your vagina at night for 3 to 7 nights. Many of these creams are available without a prescription. Your doctor can recommend the best treatment. If you would rather take a pill by mouth, you may ask your doctor for a pill called fluconazole (floo kon' na zole).

What about my partner(s)?

Usually, your sex partners do not need to be examined. However, if your infection keeps coming back, or if your partner has symptoms, your health care provider will also want to examine your partners.

How can I prevent yeast infections?

It is very important to keep a normal, healthy balance of bacteria in your vagina. You can help by:

- Keeping your genital area clean and dry;
- Wearing cotton underwear and loose fitting pants (these keep you dry by allowing air to flow through them), and avoiding douches or other feminine hygiene products.

Are there any other causes of a Vaginal Discharge?

Although most vaginal infections in women are due to bacterial vaginosis, trichomoniasis or vulvovaginal candidiasis, it is clear that there are other possible causes. These causes may include allergic and irritative factors or other STDs. Noninfectious allergic symptoms can be caused by spermicides, vaginal hygiene products, detergents or fabric softeners. Cervical infections are also often associated with abnormal vaginal discharge, but these infections can be distinguished from true vaginal infections by appropriate tests. Finally, in uninfected women, vaginal discharge may be present during ovulation and may become so heavy that it raises concern.



Information contained in this booklet is meant for informational purposes only and should not substitute the visit to your doctor nor his/her advice for your health care.

Accuracy of the content is current to the date of printing.


January 2009



DR. RAYMOND MANSOOR

*B Sc , MB BS , DM (O&G), FACOG
OBSTETRICIAN & GYNAECOLOGIST*

TANNER ST. & CORN ALLEY
P.O. BOX W1361
ST. JOHN'S
ANTIGUA
TEL: (268) 463 2232/3
info@mansoormedical.org
www.mansoormedical.org



What is a Yeast Infection? *Candidal Vaginitis or Vaginal Candidiasis*

A yeast infection is the second most common cause of vaginal discharge for women in the United States.

What causes a yeast infection?

Most of the time this infection is caused by a tiny fungus called “Candida albicans,” “Candida,” or yeast. It is normal to have a small amount of Candida in your vagina. Most people also have Candida in their mouth and lower intestinal tract. This fungus usually causes no symptoms.

A healthy vagina has a balance between healthy bacteria and unhealthy organisms. An infection occurs when something happens to allow the Candida fungus to outnumber the healthy bacteria in your vagina. When a woman has too much Candida in her vagina, then we say she has a yeast infection. A yeast infection is not considered a sexually transmitted disease (STD) because you don't have to have sex to get it.

Your vagina's healthy balance may be upset by any of the following:

- Birth control pills
- Antibiotics
- Pregnancy
- Diabetes
- Deodorant tampons or perfumed douches
- Wearing tight clothes or synthetics such as nylon, span-

dex or Lycra. These fabrics may create too much warmth and moisture from sweating. Yeast grows best in warm, moist places. Wear loose cotton clothing instead.

- HIV infection

What are the signs and symptoms?

- Vaginal itching with or without irritation
- A thick, white fluid (discharge) that looks like cottage cheese and has very little odor
- Redness, swelling, and soreness of the vaginal opening and around the opening
- Burning when urinating
- Pain during intercourse



How will I know if I have a yeast infection?

To know for sure, you should visit a health care provider. He or she will give you a pelvic exam and take a sample of your discharge. A

microscope will be used to look at your discharge in the office or it

will be sent to a lab for test-

ing. There are other types of vaginal infections with symptoms similar to a yeast infection but

they will not respond to medicine for a yeast infection. This is why it is important to visit a health care provider so you can be sure what infection you have.

When can I have sex again?

It is best to wait to have any kind of sex — oral, vaginal or anal — until you have taken all of your medicine and all of your symptoms have gone away.

Can I get this infection again?

Yes. It's not uncommon to have a second yeast infection, although it's rare to have repeated infections.



A yeast infection is the second most common cause of vaginal discharge for women in the United States.